



ELSEVIER

Available online at www.sciencedirect.com ScienceDirect

Systematic and Applied Microbiology 32 (2009) 571–582

SYSTEMATIC AND
APPLIED MICROBIOLOGYwww.elsevier.de/syapm

Water column anammox and denitrification in a temperate permanently stratified lake (Lake Rassnitzer, Germany)

M. Robert Hamersley^{a,*}, Dagmar Woebken^a, Bertram Boehrer^b, Martin Schultze^b, Gaute Lavik^a, Marcel M.M. Kuypers^a

^aMax Planck Institute for Marine Microbiology (MPIMM), Bremen, Germany

^bUFZ Helmholtz Centre for Environmental Research, Magdeburg, Germany

Received 23 April 2009

Abstract

We studied microbial N₂ production via anammox and denitrification in the anoxic water column of a restored mining pit lake in Germany over an annual cycle. We obtained high-resolution hydrochemical profiles using a continuous pumping sampler. Lake Rassnitzer is permanently stratified at ca. 29 m depth, entraining anoxic water below a saline density gradient. Mixed-layer nitrate concentrations averaged ca. 200 μmol L⁻¹, but decreased to zero in the anoxic bottom waters. In contrast, ammonium was <5 μmol L⁻¹ in the mixed layer but increased in the anoxic waters to ca. 600 μmol L⁻¹ near the sediments. In January and October, ¹⁵N tracer measurements detected anammox activity (maximum 504 nmol N₂ L⁻¹ d⁻¹ in ¹⁵NH₄⁺-amended incubations), but no denitrification. In contrast, in May, N₂ production was dominated by denitrification (maximum 74 nmol N₂ L⁻¹ d⁻¹). Anammox activity in May was significantly lower than in October, as characterized by anammox rates (maximum 6 vs. 16 nmol N₂ L⁻¹ d⁻¹ in incubations with ¹⁵NO₃⁻), as well as relative and absolute anammox bacterial cell abundances (0.56% vs. 0.98% of all bacteria, and 2.7 × 10⁴ vs. 5.2 × 10⁴ anammox cells mL⁻¹, respectively) (quantified by catalyzed reporter deposition-fluorescence *in situ* hybridization (CARD-FISH) with anammox bacteria-specific probes). Anammox bacterial diversity was investigated with anammox bacteria-specific 16S rRNA gene clone libraries. The majority of anammox bacterial sequences were related to the widespread *Candidatus* Scalindua sorokinii/brodiae cluster. However, we also found sequences related to *Candidatus* S. wagneri and *Candidatus* Brocadia fulgida, which suggests a high anammox bacterial diversity in this lake comparable with estuarine sediments.

© 2009 Elsevier GmbH. All rights reserved.

Keywords: Anammox; Denitrification; Water column; Meromictic; Lake; CARD-FISH

Introduction

Annual global anthropogenic fixation of nitrogen (N) into bioavailable forms represents an increase over natural N fixation of >100% [15]. This additional N has contributed to the eutrophication of both freshwater and marine environments, characterized by excess primary production and consequent water column

Abbreviations: CARD, catalyzed reporter deposition; CTD, conductivity, Temperature, and depth device; FISH, fluorescence *in situ* hybridization; PCR, polymerase chain reaction.

*Corresponding author at: Soka University of America, 1 University Dr., Aliso Viejo, California, USA. Tel.: +1 949 480 4409; fax: +1 949 480 4263.

E-mail address: rhamersley@soka.edu (M. Robert Hamersley).

anoxia, which imperils aquatic metazoan ecosystems. The microbial transformation of dissolved inorganic N to gaseous forms is a pivotal sink that balances the flux of N into the biosphere via N fixation and that can ameliorate the effects of excess anthropogenic N (reviewed in [34]). Denitrification, the anaerobic heterotrophic reduction of nitrate (NO_3^-) to dinitrogen gas (N_2), was long thought to be the only significant metabolic pathway of N_2 production, but classical heterotrophic denitrification now shares its position with chemolithotrophic denitrification (i.e. with sulfide or iron) [7], as well as the more recently discovered anaerobic ammonium oxidation (anammox) process [43]. Anammox, the anaerobic equimolar oxidation of ammonium (NH_4^+) with nitrite (NO_2^-) to generate N_2 , is the chemolithotrophic metabolic pathway of a diverse, but apparently monophyletic clade of planctomycete bacteria [42]. Both denitrification and anammox have been shown to play important roles in coastal sediments, but evidence accumulating since 2003 suggests that anammox may in fact be the dominant N-loss process in suboxic marine water columns [18]. The identities of the metabolic processes responsible for N_2 production have significant implications for the N cycle, as well as for carbon (C) and oxygen (O_2) budgets, particularly in water columns [28].

Although anammox has been shown to play a significant role in marine systems, less attention has been paid to freshwater and lacustrine (lake) systems. Denitrification is thought to play a significant role in lacustrine N cycling (reviewed in [34]); however, earlier methods could not differentiate between denitrification and anammox. More recently, N_2 production via anammox activity has been detected in low-salinity estuarine sediments, where its contribution to total N_2 flux ranged up to 22% [11,35,46].

As anthropogenic eutrophication increases nutrient loading to fresh waters, and these nutrients increase the frequency and duration of water column anoxia in lakes, there is an increasing need for an improved understanding of denitrification and anammox [47]. However, little is known about the importance and distribution of anammox in lakes, particularly within the water column. Early work detected the production of labeled N_2 from a $^{15}\text{NO}_3^-$ tracer in subarctic lake water [14]. Other lake studies using mass balance, isotope fractionation, or acetylene inhibition approaches also suggested N_2 production that was identified as water column denitrification [3,29,49]. Recently, anammox activity was detected in the anoxic water column of a deep, permanently stratified tropical lake (Lake Tanganyika), where on one date, anammox was estimated to contribute 13% of the total N_2 production, with the balance accounted for by denitrification [40].

Although the importance of anammox in a variety of environments has been demonstrated, little is yet known

about the factors that control the spatial and temporal distribution of denitrification *vis-à-vis* anammox. A number of studies have examined the role of organic carbon and NO_3^- availability on anammox in sediments, but have been somewhat inconclusive (reviewed in [13]). The factors controlling the relative contribution of anammox and denitrification to N_2 production in water columns are even less well understood. In addition, the significance of denitrification to water column N_2 production remains unresolved because the few published instances of significant directly measured water column denitrification (e.g. [12,19,40]) have relied on long *in vitro* incubations of 48 h or more.

As in marine systems, the anoxic conditions necessary to support lacustrine water column denitrification and/or anammox arise when density stratification of the water column prevents atmospheric replenishment of dissolved O_2 consumed by respiration. Water column stratification can be maintained by thermal density gradients, which may be ephemeral or seasonal, or permanent, as in deep lakes. Meromixis (permanent stratification) may also be maintained by salinity-related density gradients (haloclines). Deep-sea brine pools result from the hydrogeothermal solution of mineral deposits. In coastal lakes, estuaries, and fjords, haloclines form by the intrusion of seawater. Solution lakes develop haloclines via the dissolution of minerals by groundwater. The resulting halostratification can lead to the formation of very steep oxygen and chemical gradients (chemoclines), which support thin “lenses” of microorganisms adapted to particular chemical microenvironments [10]. The entrainment of particular microbial communities may be enhanced by the attendant steep density gradients (pycnoclines), which trap particles of different densities at specific depths [45]. Special sampling techniques are required to study microbial metabolism in such steep halostabilized chemoclines [10,22].

In the present report, we studied physical and chemical profiles and N_2 -production mechanisms within the monimolimnion (unmixed bottom waters) of a halostratified meromictic solution lake in northern Germany on three occasions distributed over an annual cycle. Lake Rassnitzer is a restored mining pit lake. Highly saline groundwater seepage had filled the bottom 6 m of the lake before it was filled in 1998 with diverted river water. To study anammox and denitrification in the water column of the lake, we measured hydrochemical properties in profiles at the high vertical resolution appropriate for studying steep chemoclines, and used short incubations with isotopic tracers to measure the metabolic activities of anammox and denitrification. We also looked for molecular markers of anammox bacteria in order to understand their distribution, abundance, and diversity within the lake.

Materials and methods

Study site

Lake Rassnitzer (formerly known as Merseburg-Ost 1b) is a restored mining pit lake located near Leipzig, Germany. The physical limnology of the lake and the stability of its halocline have been previously characterized [4,5,16,48]. During four field campaigns on 27 January, 19 May, 14 July, and 18 October 2005, we characterized the hydrochemical structure of the water column. Further, in January, May, and October, we performed incubations of halocline waters with the isotopically labeled tracers $^{15}\text{NH}_4^+$ and $^{15}\text{NO}_3^-$ to determine the pathways of N_2 formation (denitrification and/or anammox). We also collected water samples for detection of anammox bacteria-related 16S ribosomal ribonucleic acid (rRNA) gene sequences and for catalyzed reporter deposition-fluorescence *in situ* hybridization (CARD-FISH). All sampling and instrument casts were made from a station over the deepest portion of the lake (36 m).

Water column profiles

We obtained profiles of salinity, temperature, and oxygen with a conductivity, temperature, and depth (CTD) device (YSI 600XLM, Yellow Springs, Ohio), equipped with a Clark-type pulsed polarographic electrode (13 μm high-sensitivity membrane, 6 s response time to a 75% increase in O_2 saturation). O_2 profiles were measured in upcasts at a vertical velocity of 1 m min^{-1} . pH, turbidity, and chlorophyll fluorescence were measured with a second multiparameter CTD device (Idronaut Ocean Seven 316, Brugherio, Italy). Water samples for high-resolution nutrient analyses and ^{15}N incubations were obtained by a custom pump-CTD system employing a progressive cavity pump to minimize disruption of *in situ* particulates (Geotech Keck SP, Denver, Colorado). Water samples were collected continuously while the pump was lowered at a constant rate; sample depths were determined from the known pumping rate and were intercalibrated with the CTD data using conductivity measurements. Dye tracer studies demonstrated that “smearing” as the sample was pumped to the surface limited vertical resolution to $\pm 0.15\text{ m}$ (data not shown). In May and October, water samples for nutrient analysis were immediately acidified ($\text{pH}=2$) with HCl to prevent the formation of Fe oxide particulates upon aeration of anoxic waters; all samples were held at 4 °C during transport to our laboratory (3 h), where they were filtered (0.2 μm) and frozen ($-20\text{ }^\circ\text{C}$) for later analysis for NO_3^- , NO_2^- , NH_4^+ , and total Fe (detection limits 0.1, 0.1, 0.3, and 0.1 $\mu\text{mol L}^{-1}$, respectively). Nitrate and NO_2^- analysis were by

reduction to nitric oxide followed by chemiluminescence detection [6]. Ammonium analysis was by the colorimetric indophenol method [17]. Total Fe was determined by the ferrozine method [17]. Mixed-layer PO_4^{3-} concentrations were determined by the molybdate method [17]. Monimolimnetic sulfide (S^{2-}) was determined in May only by the methylene blue method [17]. All sampling casts within and across sampling dates were intercalibrated with reference to the 18.0 psu (practical salinity units) salinity isopleth, which is known to be stable over the annual cycle of our study [4].

^{15}N incubations and analysis

^{15}N -labeling experiments were performed in January, May, and October 2005. Water was collected by pump-CTD from 4 to 6 depths beginning within the oxycline and extending 2–4 m into the monimolimnion. Lake water was collected in quadruply-flushed serum bottles (to eliminate atmospheric contamination), which were then stoppered, sealed, and stored at *in situ* temperatures for transport back to the laboratory (3 h). The ^{15}N incubations were begun immediately upon reaching the laboratory. The experimental method was slightly modified from that previously published by Dalsgaard et al. [12]. Briefly, 250 mL of lake water were flushed with He for 15 min to remove background N_2 and amended with $^{15}\text{NH}_4^+$, $^{15}\text{NH}_4^+ + ^{14}\text{NO}_2^-$, or $^{15}\text{NO}_3^-$ (Campro Scientific, Berlin, Germany). Each of the substrates ($^{15}\text{NH}_4^+$, $^{14}\text{NO}_2^-$, and $^{15}\text{NO}_3^-$) was added to a final concentration of 8.3 $\mu\text{mol L}^{-1}$ in January and October and to 20.8 $\mu\text{mol L}^{-1}$ in May, with the exception of October, when $^{15}\text{NH}_4^+$ was added to a final concentration of 33.3 $\mu\text{mol L}^{-1}$.

Isotopically labeled lake water was transferred into 12-mL Exetainers (Labco, High Wycombe, UK) and incubated up to 24 h in the dark at *in situ* temperatures. The Exetainers were destructively sampled at ca. 0, 8, 16, and 24 h using HgCl_2 to stop biological activity. $^{14}\text{N}^{15}\text{N}:^{14}\text{N}^{14}\text{N}$ and $^{15}\text{N}^{15}\text{N}:^{14}\text{N}^{14}\text{N}$ ratios of He-equilibrated headspaces were determined by gas chromatography/isotope ratio mass spectrometry (Fisons VG Optima, UK). Estimates of total N_2 production ($^{14}\text{N}^{14}\text{N}$, $^{14}\text{N}^{15}\text{N}$, and $^{15}\text{N}^{15}\text{N}$) from anammox and/or denitrification were derived from the isotope pairing equations [43].

Catalyzed reporter deposition-fluorescence *in situ* hybridization (CARD-FISH) of anammox bacteria

Bacterioplankton samples were obtained with our custom pump-CTD system (above). Water samples were fixed with particle-free paraformaldehyde solution (final concentration, 1% v/v) for 1 h at room temperature or

overnight at 4 °C. Aliquots of 10–30 mL were vacuum-filtered onto membrane filters (GTTP, 47 mm diam., 0.22 µm pores, Millipore, Schwalbach, Germany) and stored at –20 °C until further processing. The CARD-FISH procedure was as described in Pernthaler et al. [33] and Woebken et al. [50]. The probe AMX-368 [38] as well as a mixture of probes AMX-820 [37], BS-820 [26], and BS-820C [18] were chosen to target the entire group of known anammox bacteria. After CARD-FISH, filter sections were treated with 4,6-diamidino-2-phenylindole (DAPI, a non-specific DNA stain) and subsequently evaluated by microscopy (Carl Zeiss Axioplan, Jena, Germany). We counted a minimum of 2000 DAPI-stained cells per filter section (equivalent to approximately 20 fields of view). The relative abundance of anammox bacteria labeled with the probe mix (above) was determined as a percentage of the DAPI-stained cells. Quantification of the labeled cells was performed manually on 3 replicates per water depth and errors are standard deviations.

Construction of 16S rRNA gene clone libraries

Clone libraries of Lake Rassnitzer anammox bacteria from January 2005 (30 m depth) were constructed following the protocol for the filter polymerase chain reaction (PCR) [24] using filters prepared as described above. For the PCR reaction, cells on ca. 1 mm² filter sections were lysed with 3 freeze–thaw cycles (–80 and 42 °C) in sterile PCR-grade water. One planctomycete-specific clone library (primers PLA46F [31] and 1392R [41]) and one anammox bacteria-specific clone library (primers AMX368 and 1392R) were constructed (32 PCR cycles; annealing temperatures 58 and 56 °C, respectively) following standard PCR protocols including an optimized reduction of PCR cycles to minimize PCR bias.

Three clone libraries were also constructed from DNA isolated from an October 2005 Lake Rassnitzer water sample. For DNA extraction, 450 mL of untreated water were filtered as described above. DNA was isolated from the filter from the 29-m depth according to Zhou et al. [53]. For primer pair PLA46F/1392R, four replicates were conducted with 28 PCR cycles and 58 °C annealing temperature. Three replicate PCR reactions each were conducted for primer pairs AMX368F/1392R and PLA46F/AMX820R, with 35 PCR cycles and 56 °C annealing temperature. For the latter two pairs, a preparative gel was necessary since multiple bands were observed on agarose gel. PCR reactions were purified using the QIAquick Gel Extraction Kit (QIAGEN, Hilden, Germany), prior to cloning with TOPO TA Cloning kits for sequencing with vector pCR4 (Invitrogen, Karlsruhe, Germany) according to the manufacturer's instructions. Clones were screened by PCR

for inserts of correct size and the PCR products were sequenced with the Big Dye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Darmstadt, Germany).

Phylogenetic analysis

16S rRNA sequences of sequenced PCR products were edited with the Sequencing Analysis software (Applied Biosystems) and analysed for their closest relatives using the ARB 16S rRNA gene database (version 2004 with continuous updates for anammox bacterial-related sequences) or GenBank using the BLAST program [1]. Plasmids were isolated from clones containing anammox bacterial-related 16S rRNA genes with the Montage Plasmid Miniprep96 Kit (Millipore) and inserts were completely sequenced with vector primers M13F and M13R. Sequences of about 800 and 1050 base pair length, respectively, were assembled from individual reads with the Sequencher v4.5 software (Gene Codes Corporation, Ann Arbor, USA).

The 16S rRNA gene sequences were investigated for the presence of chimeric sequences by using the CHIMERA_CHECK program from the RDP II [9] and imported into the ARB 16S rRNA gene database. Phylogenetic analyses of the 16S rRNA sequences were conducted using distance matrix, maximum parsimony, and maximum likelihood algorithms in ARB [30] with and without 50% variability filters, followed by construction of a consensus tree. Sequence identities were calculated using the similarity function of the distance matrix algorithm in ARB. Anammox bacteria-related sequences are published in GenBank under accession numbers FJ830380–FJ830386.

Results

In order to understand the role of anammox and denitrification in water column N₂ production in Lake Rassnitzer, we developed profiles of hydrochemical properties through the anoxic bottom waters, incubated samples of anoxic water with ¹⁵N-labeled substrates, measuring the production of ¹⁵N-labeled N₂, and investigated the abundance and diversity of anammox cells using molecular approaches.

Hydrochemical structure of the water column

Summer solar heating of Lake Rassnitzer surface water caused seasonal thermal stratification at ca. 10 m depth (Fig. 1A). However, the lake was permanently stratified by dense saline water which formed a steep halocline beginning at ca. 29 m (Figs. 1B and G). Lake water salinity ranged from 3 psu in the mixed upper




Fig. 1. Water column profiles of temperature, salinity, $[O_2]$, turbidity, and chl *a* fluorescence (top set of 10 panels: A–J), and pH, $[NO_3^-]$, $[NH_4^+]$, and total $[Fe]$ (lower set of 8 panels: K–R) on four sampling dates (27 January, 19 May, 14 July, and 18 October) at Lake Rassnitzer. Upper row of panels in each set (panels A–E and K–N) shows the entire water column. The oxycline region (shaded gray) is shown expanded in the lower row of panels (F–J and O–R).

layer, or mixolimnion, to 23 psu at the lake bottom. The resulting abrupt change in water density entrained an unmixed deep layer, or monimolimnion, bounded at its upper reach by a very steep oxycline, where dissolved O_2 concentrations fell from near-saturation in the mixolimnion to near-zero over a vertical distance of ca. 30 cm (Figs. 1C and H). Turbidity was constant throughout the mixolimnion, but a lens of high turbidity was found in May and July within the oxycline (Figs. 1D and I). Peaks in chlorophyll fluorescence likely associated with living phytoplankton were observed within the mixolimnion in May and July, but the high

monimolimnetic fluorescence likely indicated an accumulation of humic material and/or pheophytin [8] (Figs. 1E and J). The pH of lake waters ranged from ca. 7 to 8, with the highest values found in summer surface waters, and the lowest in monimolimnetic waters (Figs. 1K and O).

High-resolution water sampling with our pump-CTD allowed nutrient profiles to be resolved to ± 0.15 m. Nitrate concentrations were ca. $200 \mu\text{mol L}^{-1}$ in the mixolimnion, but decreased to near-zero at the top of the monimolimnion (Figs. 1L and P). Nitrite was always below the detection limit of $0.2 \mu\text{mol L}^{-1}$ (data not

shown). Ammonium concentrations were ca. $600 \mu\text{mol L}^{-1}$ near the sediments, but declined gradually through the monimolimnion to reach $<5 \mu\text{mol L}^{-1}$ within the mixolimnion (Figs. 1M and Q). Total Fe concentrations showed the same pattern, ranging from up to $1500 \mu\text{mol L}^{-1}$ near the sediments in the anoxic monimolimnion, down to $<2 \mu\text{mol L}^{-1}$ in the mixolimnion (Figs. 1N and R). Bottom water concentrations of both NH_4^+ and Fe rose over the course of the study year (Figs. 1Q and R). Phosphate within the mixolimnion was always undetectable ($<0.05 \mu\text{mol L}^{-1}$; data not shown). Although sulfide (H_2S) was detectable by its characteristic odor below 32 m in January (pers. obs.), H_2S was undetectable ($<1 \mu\text{mol L}^{-1}$) when it was measured in May (data not shown), and no detectable smell was present in monimolimnetic samples in July or October.

The zones of O_2 and NO_3^- consumption began at ca. 28.5 m depth, and averaged ca. 0.5 and 1 m in thickness, respectively, before O_2 and NO_3^- became undetectable (Figs. 1H and P). The oxycline grew thinner (<0.3 m) in May and July, as a result of high specific oxygen consumption rates in those months (14.0 and $5.8 \mu\text{mol O}_2 \text{L}^{-1} \text{d}^{-1}$, respectively, compared to 1.3 and $0.6 \mu\text{mol O}_2 \text{L}^{-1} \text{d}^{-1}$ in January and October). Seasonal changes in specific nitrate consumption rates were less dramatic, but also were highest in the spring and summer (0.26 and $0.39 \mu\text{mol N L}^{-1} \text{d}^{-1}$ in May and July, compared to 0.21 and $0.20 \mu\text{mol N L}^{-1} \text{d}^{-1}$ in January and October).

^{15}N -labeling incubations

We anaerobically incubated lake waters from below the oxycline with the isotopic tracers $^{15}\text{NO}_3^-$ or $^{15}\text{NH}_4^+$ to elucidate the contributions of the two known N_2 -producing microbial processes – anammox and denitrification – to N cycling in lake water. The production of the two labeled N_2 isotopologues – $^{15}\text{N}^{15}\text{N}$ and $^{14}\text{N}^{15}\text{N}$ – from waters amended with $^{15}\text{NH}_4^+$, $^{15}\text{NH}_4^+ + ^{14}\text{NO}_2^-$, or $^{15}\text{NO}_3^-$ is shown in Fig. 2. No $^{15}\text{N}^{15}\text{N}$ production was detected in incubations with $^{15}\text{NH}_4^+$ or $^{15}\text{NH}_4^+ + ^{14}\text{NO}_2^-$ (data omitted from Fig. 2, panels D to F for clarity).

Two distinct patterns of labeled N_2 production were seen. In January and October, $^{14}\text{N}^{15}\text{N}$ was produced from $^{15}\text{NH}_4^+$ -amended samples (Figs. 2A and C), but $^{15}\text{N}^{15}\text{N}$ production was undetectable (Figs. 2D and F), indicating that the added $^{15}\text{NH}_4^+$ combined with endogenous $^{14}\text{NO}_2^-$ to produce N_2 via anammox. There was no delay in the onset of $^{14}\text{N}^{15}\text{N}$ production, suggesting that anammox cells were metabolically active *in situ*. Amendment with $^{14}\text{NO}_2^-$ in addition to $^{15}\text{NH}_4^+$ in January and October increased anammox N_2 production over $^{15}\text{NH}_4^+$ alone at several depths (Figs. 2A and C). The absence of $^{15}\text{N}^{15}\text{N}$ production




Fig. 2. ^{15}N -labeled N_2 production during 24-h incubations of anoxic Lake Rassnitzer water with ^{15}N -labeled inorganic substrates (see legend) (note different x -axis scale in panel E). All rates are *in vitro* ^{15}N - N_2 production, not measures of anammox or denitrification rates. Top row of panels (A–C) shows singly labeled $^{14}\text{N}^{15}\text{N}$ production. Bottom row of panels (D–F) shows doubly labeled $^{15}\text{N}^{15}\text{N}$ production. No $^{15}\text{N}^{15}\text{N}$ production was detected in incubations with $^{15}\text{NH}_4^+$ or $^{15}\text{NH}_4^+ + ^{14}\text{NO}_2^-$ (data omitted for clarity). In January and October, the production of $^{14}\text{N}^{15}\text{N}$ alone in incubations with $^{15}\text{NH}_4^+$ or $^{15}\text{NO}_3^-$ indicates anammox activity, but no denitrification. In May, the production of $^{15}\text{N}^{15}\text{N}$ in incubations with $^{15}\text{NO}_3^-$ indicates denitrification activity. For estimations of rates of anammox and denitrification N_2 production derived from the isotope pairing equations [43] (see Table 1). Anammox bacterial abundances were determined by CARD-FISH in May and October. Anammox bacteria were detected, but not enumerated in January.

in $^{15}\text{NO}_3^-$ -amended incubations – despite the substantial degree of ^{15}N labeling of the *in situ* NO_3^- pool ($^{15}\text{N}/^{14}\text{N}$ ratio $F_N > 0.15$) – shows that denitrification was not a significant contributor to N_2 production during these months (Figs. 2D and F). Since denitrification produces N_2 from two N atoms that originate from NO_3^- , incubations with a high degree of ^{15}N labeling of the NO_3^- pool would be expected to produce significant amounts of $^{15}\text{N}^{15}\text{N}$ if denitrification were active [13,27].

A contrasting pattern was observed in May, when no labeled N_2 was produced in $^{15}\text{NH}_4^+$ -amended

incubations (Figs. 2B and E). However, $^{15}\text{NO}_3^-$ -amended incubations produced primarily doubly labeled $^{15}\text{N}^{15}\text{N}$, accompanied by a smaller amount of $^{14}\text{N}^{15}\text{N}$, consistent with N_2 production via denitrification (Figs. 2B and E) (see above). Estimates of the rates of total N_2 production ($^{14}\text{N} + ^{15}\text{N}$) by anammox and denitrification were made using the isotope pairing equations [43] (Table 1). Denitrification accounted for all of the labeled N_2 flux at the upper two depths, but anammox was estimated to have contributed nearly 20% and 50% of the total N_2 flux at the lower two depths, respectively (Table 1). No anammox was detected in incubations with $^{15}\text{NH}_4^+$ at these depths, possibly because $^{14}\text{N}^{15}\text{N}$ production was below detection limit in these incubations where $^{15}\text{NH}_4^+$ labeling was low ($F_N < 0.13$). Although the oxidized N substrate for anammox proper is NO_2^- rather than NO_3^- , anammox N_2 production was detected from labeled $^{15}\text{NO}_3^-$ in May and October, indicating that, as in marine water columns, anammox likely was coupled to nitrate reduction to nitrite (Figs. 2B and C), e.g. [18,27,28]. In contrast, in January, no labeled N_2 was produced in incubations with $^{15}\text{NO}_3^-$. Both anammox and denitrification activity began immediately below the oxycline, in a zone where both NO_3^- and NH_4^+ were found simultaneously (Figs. 1 and 2). However, activity was also found in deeper layers where NO_3^- was undetectable.

Abundance and phylogeny of anammox bacteria

We were not able to use standard fluorochrome-labeled fluorescence *in situ* hybridization (FISH) probes because of high sample background fluorescence. Instead, we conducted catalyzed reporter deposition (CARD)-FISH with horseradish peroxidase-labeled probes. Anammox bacteria were undetectable in aerobic mixolimnetic (mixed layer) waters, but were found in a ca. 2-m-thick stratum near the top of the anoxic hypolimnion during all of the three sampling dates for which N_2 production was also measured (January, May, and October) (Table 1). Although we were able to visually detect anammox bacteria via CARD-FISH in January at the 29.8- and 30.7-m depths, we were unable to enumerate them because of the high density of iron oxide particulates that formed when the anoxic waters became aerated during filtration. May and October water samples were acidified to prevent the formation of particulates, which permitted CARD-FISH enumeration of anammox bacteria (Figs. 2B and C). Anammox bacteria were detected between ca. 29 and 31 m water depth on both of these dates. The abundance of anammox bacteria relative to total bacterial numbers (DAPI count) within the N_2 production zone was significantly lower in May (mean $0.56 \pm 0.06\%$) than in October (mean $0.98 \pm 0.03\%$) (one-tailed *t*-test, d.f. = 17, $p = 8 \times 10^{-6}$). Anammox cell numbers in the

Table 1. Estimated N_2 production (total $^{14}\text{N} + ^{15}\text{N}$) by anammox and denitrification calculated with the isotope pairing equations [43] in incubations of Lake Rassnitzer water with three different combinations of ^{15}N -labeled inorganic N substrates.

Date	Depth (m)	N_2 production rates (total $^{14}\text{N} + ^{15}\text{N}$) ($\text{nmol N}_2 \text{ L}^{-1} \text{ d}^{-1}$)			
		Anammox			Denitrification
		$^{15}\text{NH}_4^+$	$^{15}\text{NH}_4^+ + ^{14}\text{NO}_2^-$	$^{15}\text{NO}_3^-$	$^{15}\text{NO}_3^-$
27 Jan 2005	28.8	50 (10)	10 (2)	nd	nd
	29.8 ^a	nd	220 (30)	nd	nd
	30.7 ^a	540 (50)	2600 (600)	nd	nd
	31.6	nd	nd	nd	nd
	33.5	nd	nd	nd	nd
19 May 2005	28.7	nd	nd	nd	4.7 (0.3)
	29.3 ^b	nd	nd	nd	74 (5)
	29.8 ^b	nd	nd	6 (1)	23 (2)
	30.5 ^b	nd	nd	4 (1)	4 (2)
18 Oct 2005	28.3	nd	8 (1)	nd	nd
	28.7 ^a	nd	nd	nd	nd
	29.0 ^b	nd	10 (2)	16 (1)	nd
	29.3 ^b	nd	nd	1.8 (0.2)	nd
	29.6 ^b	14.2 (0.4)	37 (8)	13.2 (0.2)	nd
	30.2 ^b	nd	nd	nd	nd

nd indicates non-detection of N_2 production. Numbers in parentheses are standard errors (from linear regressions).

^aAnammox cells detected but could not be quantified.

^bAnammox cells detected and quantified at this depth with CARD-FISH.




Fig. 3. Phylogenetic tree showing the affiliation of representative Lake Rassnitzer anammox bacterial 16S rRNA gene sequences. The consensus tree was built based on maximum likelihood, neighbor joining, and maximum parsimony trees, calculated with and without 50% position variability filters using other cultured and uncultured planctomycete-sequences as an outgroup. The bar represents 10% estimated sequence divergence. Anammox bacterial sequences from Lake Rassnitzer are depicted in bold and all sequence descriptions are followed by their GenBank accession number and sequence length (in base pairs) in parentheses.

N_2 -production zone were also significantly lower in May (mean $2.7 \pm 0.3 \times 10^4 \text{ mL}^{-1}$) than in October (mean $5.2 \pm 0.2 \times 10^4 \text{ mL}^{-1}$) (one-tailed t -test, d.f. = 17, $p = 2 \times 10^{-6}$) (Figs. 2B and C). Previous work has shown that CARD-FISH may underestimate the abundance of anammox bacteria [50], so these numbers should be considered minima. The significantly lower absolute and relative anammox bacterial cell numbers in May compared to October are consistent with the lower anammox N_2 production rate in that month (Table 1). Cell-specific anammox rates in May ranged from 0.4 to $0.5 \text{ fmol N d}^{-1}$, whereas in October, the range was 0.08 – $0.6 \text{ fmol N d}^{-1}$.

Clone libraries from January and October 2005 constructed with the planctomycete-specific primer Pla46F and the universal primer 1392R did not result

in any sequences related to anammox bacteria. The anammox bacteria-specific forward primer Amx-368F with 1392R also failed to produce anammox bacteria-related sequences in October samples. However, the primer combinations Amx368F/1392R and Pla46F/Amx820R retrieved 6 anammox bacteria-related sequences from the January sample and 58 from the October sample, respectively.

The majority (61 of 64) of the anammox bacterial 16S rRNA gene sequences from Lake Rassnitzer formed one cluster (based on 99% sequence identity) and were closely related to *Candidatus Scalindua brodae* [38] (98.8–99.1% sequence identity). These sequences are represented in Fig. 3 by the clones clone_32u, clone_1o, MB1_3H, and MB1_4B. The remaining 3 Lake Rassnitzer sequences (of the 64) clustered

independently. Clone_23u showed 95.1% sequence identity with *Candidatus S. wagneri* [38]. However, it was most closely related to two partial sequences from sediments of the Yodo River Estuary, Japan (AB300483 and AB300470, 98.9% sequence identity) [2] and two sequences from sediments of the Cape Fear River Estuary, North Carolina, USA (FJ490127 and FJ490132, 98.7% sequence identity) [11]. Clone_29u and clone_37o were related to *Candidatus Brocadia fulgida* (94.1% and 93.2% sequence identity, respectively), an anammox bacterium enriched from an anammox wastewater treatment plant in Rotterdam, The Netherlands [23]. The closest relatives of clone_29u were a sequence from freshwater sediments of the Xinyi River (DQ198371, 99.8% sequence identity) [52] and sequences from Yodo River Estuary sediment (AB300485 and AB300480, 99.3% sequence identity) [2]. Clone_37o was most closely related to a sequence from Cape Fear River Estuary sediments (FJ490119, 99.6% sequence identity) [11].

Discussion

Seasonal variation of anammox and denitrification in the Lake Rassnitzer water column

Our measurements of N_2 production patterns in the anoxic monimolimnion of Lake Rassnitzer showed an unexpected seasonal shift in the dominant process responsible for water column N_2 production – from anammox in the fall and winter to denitrification in the spring (Fig. 2, Table 1). The reasons for this shift were not clear. Water temperatures at the depths where anammox and denitrification were measured are quite stable throughout the year ($\pm 1^\circ\text{C}$), as were the concentrations of NO_3^- and NH_4^+ (Fig. 1). Although anammox activity was detected in all sampling dates (Fig. 2), anammox N_2 production rates estimated from our isotope tracer experiments were lowest in May, the only date when denitrification was detected (Table 1). The increase in anammox rates from May to October (from a maximum of 6–16 $\text{nmol N}_2 \text{L}^{-1} \text{d}^{-1}$) was accompanied by a concomitant and significant increase in relative and absolute anammox cell abundances (from 0.56% to 0.98% of DAPI bacteria, and from 2.7×10^4 to 5.2×10^4 anammox cells mL^{-1} , respectively). Cell-specific anammox rates remained relatively unchanged (maxima 0.5 and 0.6 fmol N d^{-1}), suggesting that the increase in anammox activity in October was due to the growth of anammox bacteria. Thus the early summer appears to be a relatively unfavorable season for anammox bacterial growth.

Evidence from marine sediments suggests that the contribution of anammox to total N_2 production in

sediments increases as indices of organic loading decrease (e.g. chlorophyll *a*, decreasing water depth, O_2 consumption) (reviewed in [13]). This trend has been proposed to result from an increase in competition for NO_2^- by heterotrophic denitrifiers when sufficient reductant is present. Although there is insufficient evidence to suggest that a similar relationship between organic loading and denitrification exists in the water column, we observed a sharp summer increase in O_2 consumption within the oxycline from $< 1.3 \mu\text{mol O}_2 \text{L}^{-1} \text{d}^{-1}$ in January and October (when anammox alone was detected) to $14.0 \mu\text{mol O}_2 \text{L}^{-1} \text{d}^{-1}$ in May (when denitrification dominated), suggestive of increased organic C oxidation in May. Water column denitrification in May might also have been driven by the availability of reductants other than organic C, such as reduced Fe or sulfide. A similar, but *interannual* shift from denitrification to anammox was observed in the Baltic Sea, where it was attributed to the removal of sulfide substrate for denitrification by an influx of oxygenated water [19]. Although sulfide was not generally available as a reductant for chemoautotrophic denitrification in Lake Rassnitzer, reduced Fe was present in very high concentrations all year (Fig. 1), and chemolithotrophic iron-oxidizing bacteria are known from lake sediments [20]. However, the identity of the reductant for Lake Rassnitzer denitrification as well as the controls on the relative importance of denitrification and anammox remain unknown.

Anammox N_2 production rates estimated from our ^{15}N incubations using the isotope pairing equations [43] (up to $540 \text{ nM N}_2 \text{d}^{-1}$ in incubations with $^{15}\text{NH}_4^+$; Table 1) ranged slightly higher than maximum rates reported earlier from marine water columns (17–430 $\text{nM N}_2 \text{d}^{-1}$ [12,18,27,44]), as well as from freshwater Lake Tanganyika (240 $\text{nM N}_2 \text{d}^{-1}$ [40]). However, most of the anammox rates we determined in Lake Rassnitzer were similar to previously reported rates. CARD-FISH-derived anammox bacterial cell numbers in Lake Rassnitzer also ranged higher ($5.2 \times 10^4 \text{ mL}^{-1}$) than previously reported maxima from marine systems ($1.3\text{--}2.2 \times 10^4 \text{ mL}^{-1}$ [19,26,27]) with the exception of the Peruvian upwelling system, where anammox cell abundances ranged up to $15 \times 10^4 \text{ mL}^{-1}$ [18]. Anammox bacteria in Lake Rassnitzer accounted for a maximum of 1.05% of total bacterial cells (DAPI), at the low end of the range reported earlier (maxima 0.75–3.1% [18,19,27]). In contrast, denitrification rates in the present study ($\leq 74 \text{ nM N}_2 \text{d}^{-1}$) were at the lower end of the range of denitrification maxima previously reported from marine and freshwater water columns (480–2700 $\text{nM N}_2 \text{d}^{-1}$ [12,19,40]).

Anammox activities observed in incubations with $^{15}\text{NH}_4^+$ vs. $^{15}\text{NO}_3^-$ were not always consistent. Anammox activity was detected in January in incubations with $^{15}\text{NH}_4^+$, but not with $^{15}\text{NO}_3^-$. Since the NO_3^- pool

in the incubations was highly labeled with ^{15}N at 30.7 m where anammox activity was high, we can only conclude that nitrate was not being reduced to nitrite to support anammox in January, unlike in May or October, or as observed in marine water columns (e.g. [18,27,28]). The opposite condition was observed in May, when anammox was observed in incubations with $^{15}\text{NO}_3^-$, but not $^{15}\text{NH}_4^+$. A possible explanation for the May observation is that the NH_4^+ pool in May was weakly labeled ($F_{\text{N}} < 0.13$) relative to the NO_3^- pool ($F_{\text{N}} > 0.89$), so low rates of anammox activity may not have resulted in the production of detectable $^{14}\text{N}^{15}\text{N}$. These differences emphasize the value of conducting incubations with multiple ^{15}N -labeled substrates.

Anammox bacterial diversity

A number of studies have investigated the diversity of anammox bacteria in marine and freshwater sediments and water columns by culture-independent approaches, using PCR with primers targeting either planctomycete or anammox bacteria-related 16S rRNA genes (e.g. [2,11,18,25–27,32,36,39,51]). However, in marine systems, only sequences related to the *Candidatus* S. brodae/sorokinii cluster have so far been recovered.

In contrast, freshwater and brackish systems appear to harbor a greater diversity of anammox bacteria. *Candidatus* Brocadia anammoxidans-related sequences were recovered from sediments of the Xinyi River in China in addition to a *Candidatus* Scalindua-related sequence (after 36 days of enrichment) [52]. A recent study of hot springs sediments and microbial mats in California and Nevada, USA, found anammox bacterial sequences related to two anammox bacterial genera, *Candidatus* Kuenenia and *Candidatus* Brocadia, but not *Candidatus* Scalindua [21]. An even higher anammox bacterial diversity was found in a Japanese estuary, where sequences clustering with *Candidatus* Scalindua, *Candidatus* Brocadia, and *Candidatus* Kuenenia were recovered from sediments of the Yodo River Estuary [2]. A similarly high diversity was found in sediments of the Cape Fear River Estuary, USA [11]. The anammox bacterial sequences recovered from Lake Rassnitzer in the present study also grouped with three distinct clusters: *Candidatus* S. sorokinii/brodae, *Candidatus* S. wagneri, and *Candidatus* B. fulgida (Fig. 3). Three of these sequences (clustering with *Candidatus* B. fulgida as well as *Candidatus* S. wagneri) were closely related to sequences from the Xinyi River, the Yodo River, and Cape Fear River estuaries (Fig. 3).

The reasons for the increased diversity in these freshwater and brackish systems are still unclear. In the Cape Fear River Estuary, an increase in anammox bacterial diversity was observed along a gradient of increasing salinity and salinity variation. The diversity of anammox

bacteria in Lake Rassnitzer is similar to that of the most haline and diverse Cape Fear River Estuary site. However, at both sites, the majority of the sequences were related to the *Candidatus* S. sorokinii/brodae cluster. The salinity of the N_2 production zone of Lake Rassnitzer ranges from ca. 6 to 13 psu, similar to the maximum salinity observed at the most haline Cape Fear River Estuary site, although the estuary site was highly temporally variable (0.4–9.9 psu) compared to Lake Rassnitzer. Dale and co-workers [11] propose that *Candidatus* Scalindua is the most halotolerant of the anammox genera and that *Candidatus* Kuenenia and *Candidatus* Brocadia are more sensitive to salinity. Our results from Lake Rassnitzer, where most of the sequences we found were related to *Candidatus* Scalindua, support this suggestion. However, we also found a sequence related to *Candidatus* Brocadia at 6.2 psu, indicating that some members of this genera may also be somewhat halotolerant. The only other known lacustrine water column 16S rRNA gene sequence (recovered from the tropical freshwater Lake Tanganyika) was most closely related to *Candidatus* S. sorokinii/brodae, but only a single sequence was recovered from this site [40].

Although anammox bacteria in lakes have been much less studied than in marine systems, our results support the possibility that freshwater and brackish environments such as lakes and estuaries may harbor a greater diversity of anammox bacteria than marine systems. Our observations of an apparent seasonal shift between anammox and denitrification in the water column of Lake Rassnitzer underline the importance of seasonal studies of N_2 production processes in lacustrine, as well as marine water columns in order to understand the relative contributions of the two processes and their impacts on suboxic nutrient balances. The continuing impact of anthropogenic nutrient inputs to lakes, and the accompanying increases in the extent of water column anoxia, are likely to increase the role of microbial N_2 production in lacustrine N balances. Our study indicates that the microbiology of N_2 production in lacustrine water columns may be more complex than previously recognized.

Acknowledgements

We thank Karsten Rahn, Uwe Kivel, and UFZ staff for logistical and field support at Lake Rassnitzer. MRH would like to especially thank K. Rahn for his kind hospitality during field campaigns. We also thank Tina Willers (University of Bremen), and Athanasios Kaffes, Sabine Lenk, and Judith Ufkes (MPIMM) for field and laboratory assistance. Funding was provided by MPIMM, UFZ, and the Wrigley Institute for Environmental Studies. We thank two anonymous reviewers for their helpful comments.

References

- [1] S.F. Altschul, T.L. Madden, A.A. Schäffer, J. Zhang, Z. Zhang, W. Miller, et al., Gapped BLAST and PSI-BLAST: a new generation of protein database search programs, *Nucleic Acids Res.* 25 (1997) 3389–3402.
- [2] T. Amano, I. Yoshinaga, K. Okada, T. Yamagishi, S. Ueda, A. Obuchi, Y. Sako, Y. Suwa, Detection of anammox activity and diversity of anammox bacteria-related 16S rRNA genes in coastal marine sediment in Japan, *Microbes Environ.* 22 (2007) 232–242.
- [3] C. Bédard, R. Knowles, Hypolimnetic O₂ consumption, denitrification, and methanogenesis in a thermally stratified lake, *Can. J. Fish Aquat. Sci.* 48 (1991) 1048–1054.
- [4] B. Boehrer, H. Heidenreich, M. Schimmele, M. Schultze, Numerical prognosis for salinity profiles of future lakes in the opencast mine Merseburg-Ost, *Int. J. Salt Lake Res.* 7 (1998) 235–260.
- [5] B. Boehrer, A. Matzinger, M. Schimmele, Similarities and differences in the annual temperature cycles of east German mining lakes, *Limnologia* 30 (2000) 271–279.
- [6] R.S. Braman, S.A. Hendrix, Nanogram nitrite and nitrate determination in environmental and biological materials by vanadium (III) reduction with chemiluminescence detection, *Anal. Chem.* 61 (1998) 2715–2718.
- [7] I. Brettar, G. Rheinheimer, Denitrification in the Central Baltic: evidence for H₂S-oxidation as motor of denitrification at the oxic–anoxic interface, *Mar. Ecol. Prog. Ser.* 77 (1991) 157–169.
- [8] R.E. Carlson, J. Shapiro, Dissolved humic substances: a major source of error in fluorometric analyses involving lake waters, *Limnol. Oceanogr.* 26 (1981) 785–790.
- [9] J.R. Cole, B. Chai, T.L. Marsh, R.J. Farris, Q. Wang, S.A. Kulam, et al., The Ribosomal Database Project (RDP-II): previewing a new autoaligner that allows regular updates and the new prokaryotic taxonomy, *Nucleic Acids Res.* 31 (2003) 442–443.
- [10] D. Daffonchio, S. Borin, T. Brusa, L. Brusetti, P.W.J.J. van der Wielen, H. Bolhuis, et al., Stratified prokaryote network in the oxic–anoxic transition of a deep-sea halocline, *Nature* 440 (2006) 203–207.
- [11] O.R. Dale, C.R. Tobias, B. Song, Biogeographical distribution of diverse anaerobic ammonium oxidizing (anammox) bacteria in Cape Fear River Estuary, *Environ. Microb.* 11 (2009) 1194–1207.
- [12] T. Dalsgaard, D.E. Canfield, J. Petersen, B. Thamdrup, J. Acuña-González, N₂ production by the anammox reaction in the anoxic water column of Golfo Dulce, Costa Rica, *Nature* 422 (2003) 606–608.
- [13] T. Dalsgaard, B. Thamdrup, D.E. Canfield, Anaerobic ammonium oxidation (anammox) in the marine environment, *Res. Microbiol.* 156 (2005) 457–464.
- [14] J.J. Goering, V.A. Dugdale, Estimates of the rates of denitrification in a subarctic lake, *Limnol. Oceanogr.* 11 (1966) 113–117.
- [15] J.N. Galloway, F.J. Dentener, D.G. Capone, E.W. Boyer, R.W. Howarth, S.P. Seitzinger, et al., Nitrogen cycles: past, present, and future, *Biogeochemistry* 70 (2004) 153–226.
- [16] H. Gräfe, B. Boehrer, N. Hoppe, S.C. Müller, P. Hauptmann, Ultrasonic in situ measurements of density, adiabatic compressibility, and stability frequency, *Limnol. Oceanogr.* 47 (2002) 1255–1260.
- [17] K. Grasshoff, M. Ehrhardt, K. Kremling, L.G. Anderson, *Methods of Seawater Analysis*, third ed., Wiley-VCH, Weinheim, 1999 596pp.
- [18] M.R. Hamersley, G. Lavik, D. Woebken, J.E. Rattray, P. Lam, E.C. Hopmans, et al., Anaerobic ammonium oxidation in the Peruvian oxygen minimum zone, *Limnol. Oceanogr.* 52 (2007) 923–933.
- [19] M. Hannig, G. Lavik, M.M.M. Kuypers, D. Woebken, W. Martens-Habbena, K. Jürgens, Shift from denitrification to anammox after inflow events in the central Baltic Sea, *Limnol. Oceanogr.* 52 (2007) 1336–1345.
- [20] S. Hauck, M. Benz, A. Brune, B. Schink, Ferrous iron oxidation by denitrifying bacteria in profundal sediments of a deep lake (Lake Constance), *FEMS Microbiol. Ecol.* 37 (2003) 127–134.
- [21] A. Jaeschke, H.J.M. Op den Camp, H. Harhangi, A. Klimiuk, E.C. Hopmans, M.S.M. Jetten, et al., 16S rRNA gene and lipid biomarker evidence for anaerobic ammonium-oxidizing bacteria (anammox) in California and Nevada hot springs, *FEMS Microbiol. Ecol.* 67 (2009) 342–350.
- [22] B.B. Jørgensen, J.G. Kuenen, Y. Cohen, Microbial transformations of sulfur compounds in a stratified lake (Solar Lake, Sinai), *Limnol. Oceanogr.* 24 (1979) 799–822.
- [23] B. Kartal, L. van Niftrik, A.O. Sliemers, M.C. Schmid, I. Schmidt, K. van de Pas-Schoonen, et al., Application, eco-physiology and biodiversity of anaerobic ammonium-oxidizing bacteria, *Rev. Environ. Sci. Biotechnol.* 3 (2004) 255–264.
- [24] D.L. Kirchman, L. Yu, B.M. Fuchs, R. Amann, Structure of bacterial communities in aquatic systems as revealed by filter PCR, *Aquat. Microb. Ecol.* 26 (2001) 13–22.
- [25] J. Kirkpatrick, B. Oakley, C. Fuchsman, S. Srinivasan, J.T. Staley, J.W. Murray, Diversity and distribution of planctomycetes and related bacteria in the suboxic zone of the Black Sea, *Appl. Environ. Microbiol.* 72 (2006) 3079–3083.
- [26] M.M.M. Kuypers, A.O. Sliemers, G. Lavik, M. Schmid, B.B. Jørgensen, J.G. Kuenen, et al., Anaerobic ammonium oxidation by anammox bacteria in the Black Sea, *Nature* 422 (2003) 608–611.
- [27] M.M.M. Kuypers, G. Lavik, D. Woebken, M. Schmid, B.M. Fuchs, R. Amann, et al., Massive nitrogen loss from the Benguela upwelling system through anaerobic ammonium oxidation, *Proc. Natl. Acad. Sci. USA* 102 (2005) 6478–6483.
- [28] P. Lam, G. Lavik, M.M. Jensen, J. van de Vossenberg, M. Schmid, D. Woebken, et al., Revising the nitrogen cycle in the Peruvian oxygen minimum zone, *Proc. Natl. Acad. Sci. USA* 106 (2009) 4752–4757.
- [29] M.F. Lehmann, P. Reichert, S.M. Bernasconi, A. Barbieri, J.A. McKenzie, Modelling nitrogen and oxygen isotope fractionation during denitrification in a lacustrine redox-transition zone, *Geochim. Cosmochim. Acta* 67 (2003) 2529–2542.

- [30] W. Ludwig, O. Strunk, R. Westram, L. Richter, H. Meier, Yadhukumar, et al., ARB: a software environment for sequence data, *Nucleic Acids Res.* 32 (2004) 1363–1371.
- [31] A. Neef, R. Amann, H. Schlesner, K.-H. Schleifer, Monitoring a widespread bacterial group: *in situ* detection of planctomycetes with 16S rRNA-targeted probes, *Microbiology* 144 (1998) 3257–3266.
- [32] C.R. Penton, A.H. Devol, J.M. Tiedje, Molecular evidence for the broad distribution of anaerobic ammonium-oxidizing bacteria in freshwater and marine sediments, *Appl. Environ. Microbiol.* 72 (2006) 6829–6832.
- [33] A. Pernthaler, J. Pernthaler, R. Amann, Fluorescence *in situ* hybridization and catalyzed reporter deposition for the identification of marine bacteria, *Appl. Environ. Microbiol.* 68 (2002) 3094–3101.
- [34] E. Piña-Ochoa, M. Álvarez-Cobelas, Denitrification in aquatic environments: a cross-system analysis, *Biogeochemistry* 81 (2006) 111–130.
- [35] J.J. Rich, O.R. Dale, B. Song, B.B. Ward, Anaerobic ammonium oxidation (anammox) in Chesapeake Bay sediments, *Microb. Ecol.* 55 (2008) 311–320.
- [36] N. Risgaard-Petersen, R.L. Meyer, M. Schmid, M.S.M. Jetten, A. Enrich-Prast, S. Rysgaard, et al., Anaerobic ammonium oxidation in an estuarine sediment, *Aquat. Microb. Ecol.* 36 (2004) 293–304.
- [37] M. Schmid, U. Twachtmann, M. Klein, M. Strous, S. Juretschko, M. Jetten, et al., Molecular evidence for genus level diversity of bacteria capable of catalyzing anaerobic ammonium oxidation, *Syst. Appl. Microbiol.* 23 (2000) 93–106.
- [38] M. Schmid, K. Walsh, R. Webb, W.I. Rijpstra, K. van de Pas-Schoonen, M.J. Verbruggen, et al., *Candidatus* “*Scalindua brodae*”, sp. nov., *Candidatus* “*Scalindua wagneri*”, sp. nov., two new species of anaerobic ammonium oxidizing Bacteria, *Syst. Appl. Microbiol.* 26 (2003) 529–538.
- [39] M.C. Schmid, N. Risgaard-Petersen, J. van de Vossenberg, M.M.M. Kuypers, G. Lavik, J. Petersen, et al., Anaerobic ammonium-oxidizing bacteria in marine environments: widespread occurrence but low diversity, *Environ. Microbiol.* 9 (2007) 1476–1484.
- [40] C.J. Schubert, E. Durisch-Kaiser, B. Wehrli, B. Thamdrup, P. Lam, M.M.M. Kuypers, Anaerobic ammonium oxidation in a tropical freshwater system (Lake Tanganyika), *Environ. Microbiol.* 8 (2006) 1857–1863.
- [41] D.A. Stahl, B. Flesher, H.R. Mansfield, L. Montgomery, Use of phylogenetically based hybridization probes for studies of ruminal microbial ecology, *Appl. Environ. Microbiol.* 54 (1988) 1079–1084.
- [42] M. Strous, J.G. Kuenen, M.S.M. Jetten, Key physiology of anaerobic ammonium oxidation, *Appl. Environ. Microbiol.* 65 (1999) 3248–3250.
- [43] B. Thamdrup, T. Dalsgaard, Production of N₂ through anaerobic ammonium oxidation coupled to nitrate reduction in marine sediments, *Appl. Environ. Microbiol.* 68 (2002) 1312–1318.
- [44] B. Thamdrup, T. Dalsgaard, M.M. Jensen, O. Ulloa, L. Farías, R. Escibano, Anaerobic ammonium oxidation in the oxygen-deficient waters off northern Chile, *Limnol. Oceanogr.* 51 (2006) 2145–2156.
- [45] J.H. Trefry, B.J. Presley, W.L. Keeney-Kennicutt, R.P. Trocine, Distribution and chemistry of manganese, iron, and suspended particulates in Orca Basin, *Geo-Mar. Lett.* 4 (1984) 125–130.
- [46] M. Trimmer, J.C. Nicholls, B. Deflandre, Anaerobic ammonium oxidation measured in sediments along the Thames Estuary, United Kingdom, *Appl. Environ. Microbiol.* 69 (2003) 6447–6454.
- [47] P.M. Vitousek, J.D. Aber, R.W. Howarth, G.E. Likens, P.A. Matson, D.W. Schindler, et al., Human alteration of the global nitrogen cycle: sources and consequences, *Ecol. Appl.* 7 (1997) 737–750.
- [48] C. von Rohden, J. Ilmberger, Tracer experiment with sulfur hexafluoride to quantify the vertical transport in a meromictic pit lake, *Aquat. Sci.* 63 (2001) 417–431.
- [49] B.B. Ward, J. Granger, M.T. Maldonado, K.L. Casciotti, S. Harris, M.L. Wells, Denitrification in the hypolimnion of permanently ice-covered Lake Bonney, Antarctica, *Aquat. Microb. Ecol.* 38 (2005) 295–307.
- [50] D. Woebken, B.M. Fuchs, M.M.M. Kuypers, R. Amann, Potential interactions of particle-associated anammox bacteria with bacterial and archaeal partners in the Namibian upwelling system, *Appl. Environ. Microbiol.* 73 (2007) 4648–4657.
- [51] D. Woebken, P. Lam, M.M.M. Kuypers, S.W.A. Naqvi, B. Kartal, M. Strous, et al., A microdiversity study of marine anammox bacteria reveals a novel *Candidatus* *Scalindua* phylotype in marine oxygen minimum zones, *Environ. Microbiol.* 10 (2008) 3106–3119.
- [52] Y. Zhang, X.-H. Ruan, H.J.M. Op den Camp, T.J.M. Smits, M.S.M. Jetten, M. Schmid, Diversity and abundance of aerobic and anaerobic ammonium-oxidizing bacteria in freshwater sediments of the Xinyi River (China), *Environ. Microbiol.* 9 (2007) 2375–2382.
- [53] J. Zhou, M.A. Bruns, J.M. Tiedje, DNA recovery from soils of diverse composition, *Appl. Environ. Microbiol.* 62 (1996) 316–322.